

QuickBayt[®] Spot Spray



QuickBayt[®] Spot Spray 3 oz in a Spray Bottle Makes Killing Flies Easier.

The proven fly-killing power of QuickBayt[®] Spot Spray is now available in a convenient 24 fl oz spray bottle. Named one of the Top Ten New Products at World Ag Expo 2008, QuickBayt Spot Spray in the smaller size is:

Convenient and Easy to Use

- Easy to apply. Just add water and spot-spray problem areas.
- For best results, make a number of spot sprays throughout the facility.
- Reusable bottle includes a built-in sprayer.
- Bottle can be refilled from economic 1 lb container.

Powerful and Long Lasting

- Starts killing flies in less than 60 seconds.¹
- Contains imidacloprid, the same active ingredient as QuickBayt Fly Bait.

- Contains two attractants that flies can't resist.
- Unless consumed prior, one application lasts up to six weeks indoors/two weeks outdoors.

Flexible

- Can be sprayed on walls, posts, beams and ceilings.
- Can be used indoors or outdoors.
- Dries virtually clear on most surfaces.



1. Nipcam Study, QuickBayt Technical Information Manual, December 2002.
© 2009 Bayer HealthCare LLC, Animal Health Division, Shawnee Mission, Kansas 66201
Bayer, the Bayer Cross and QuickBayt are registered trademarks of Bayer.



QuickBayt® Spot Spray

**For use in and around animal facilities –
designed to lure house flies to their death!**

- Add water, spot spray and watch flies pile up
- Quick acting and long lasting control
- Contains 2 attractants flies can't resist

ACTIVE INGREDIENTS:	
Imidacloprid	10.0%
Z-9-Tricosene.....	0.1%
OTHER INGREDIENTS	89.9%
TOTAL 100.0%	

EPA Reg. No. 11556-147 EPA Est. No. 33967-NJ-1

**STOP – Read the label before use.
Keep out of reach of children.**

CAUTION

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency medical treatment call 1-800-422-9874. For product information call 1-800-633-3796.	
Note To Physician: No specific antidote is available. Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

QUICK FACTS	
QuickBayt® Spot Spray	<ul style="list-style-type: none"> • A bait that is consumed by house flies. • Simple and easy to use. Just add water, mix, and spray. • Don't treat the entire area, make spot applications. • Outdoor residual efficacy up to 2 weeks unless the bait is completely consumed. • In animal facilities and protected areas residual efficacy can last up to 6 weeks unless the bait is completely consumed. • Contains Bitrex® bittering agent to help prevent ingestion by non-target animals, children and pets.
WHERE TO APPLY	<ul style="list-style-type: none"> • In and around agricultural livestock production facilities, such as: poultry houses, turkey and chicken houses, cattle, sheep and goat facilities, feed lots, dairy barns, swine confinement buildings, dog kennels, and horse stables. • In and around hobby and recreational farms and rural environments where house flies aggregate.
HOW TO APPLY	<ul style="list-style-type: none"> • Mix in water and spray on surfaces.
COVERAGE	<ul style="list-style-type: none"> • Use 24 fl oz of finished spray to treat 188 square feet of fly-resting surfaces.
RE-APPLY	<ul style="list-style-type: none"> • Re-apply as needed, but not before 7 days.
QUESTIONS ??	<ul style="list-style-type: none"> • For questions or comments, call toll-free 1-800 633-3796.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read entire label before using this product.

Information About QuickBayt® Spot Spray

This product is a bait formulated to control house flies in animal confinement areas (e.g., cattle, poultry, swine), hobby and recreational farms, and rural environments where house flies aggregate. The product can be used as a spray for treatment of surfaces, such as stable walls or waste containers. There is no need to spray the complete surface because this bait will attract pests to the treated area. Begin applications at the start of the season before fly populations have reached their peak. Best results occur when the product is applied to many locations throughout the structure or facility. Spray the product on surfaces where house flies tend to rest. House flies prefer warm, still sites for resting, so avoid cool or windy locations.

Applications of this product may leave a visible residue or discolor some surfaces. Before treating, it is advisable to test application in a small inconspicuous area.

MIXING DIRECTIONS, USE RATES AND APPLICATION

All dilutions specified on this label result in a 1.2% imidacloprid spray. Remove the trigger sprayer from the bottle. Fill the bottle with water to 24 fl oz to make a 1.2% imidacloprid dilution. Put the sprayer back on the bottle and tighten. Shake the bottle to mix the product. Spray as directed.

Directions for refill of this package:

This trigger spray bottle may be refilled and used to make additional QuickBayt Spot Spray solution. After obtaining a one pound bottle of QuickBayt Spot Spray, follow the directions below to prepare additional spray solution.

Remove the trigger sprayer from the bottle. Add 1/2 cup of QuickBayt Spot Spray to the bottle. Fill the bottle with water to 24 fl oz to make a 1.2% imidacloprid dilution. Put the sprayer back on the bottle and tighten. Shake the bottle to mix the product. Spray as directed.

HOW AND WHERE TO APPLY

Applications in food/feed establishments are not permitted.

FOR USE IN and around agricultural livestock production facilities, such as poultry houses, turkey and chicken houses, cattle, sheep and goat facilities, feedlots, dairy barns, swine confinement buildings, dog kennels, horse stables, animal confinement areas, animal holding areas, animal quarters, animal rearing facilities, livestock [pigs, cattle, sheep, goat] barns, storage areas, storage rooms, dumpsters and trash collection areas, utility rooms, washrooms, work areas such as offices and rooms in animal facilities, and hobby and recreational farms and rural environments where houseflies tend to be a nuisance or a health hazard.

Application:

Spray diluted product where house flies aggregate: on walls, around windows, on surfaces of pillars and other structures, etc. For best results, apply product in spots of 3 to 12 sq. ft. where house flies tend to rest. This package will treat 188 square feet of fly-resting surfaces.

Residual efficacy in animal facilities:

Residual efficacy in animal facilities is dependent on the sanitary conditions and fly pressure level, as well as on the quality and cleanliness of the surfaces treated.

Apply as soon as fly pressure exceeds acceptable threshold levels to avoid population build up. Re-apply as needed, typically at 4 to 6 week intervals, but no sooner than 7 days after application.

When spraying surfaces, care must be taken only to spray places out of reach of the animals. This also applies to empty pens due to be re-stocked within the next few weeks. Do not spray directly on animals, feed and drinking supplies nor on milking equipment. Do not treat surfaces that people frequently contact. Avoid contact with sprayed surfaces until dry.

Restrictions

- Do not apply bait on surfaces accessible to children or pets.
- Do not apply bait in areas where food animals can ingest product.
- Do not allow bait to come in contact with food or water.
- Do not apply to milking equipment at dairy facilities.
- Do not apply inside food processing or manufacturing facilities.
- Do not apply in classrooms or patient rooms.

For Best Results

- Use the bait as part of an overall Integrated Pest Management program.
- Use proper sanitation and manure management.
- Consult with cooperative extension service in your state for advice on fly management practices.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Refillable container. Refill this container with QuickBayt® Spot Spray (imidacloprid/Z-9-tricosene) only. Do not reuse this container for any other purpose.

Pesticide Storage:

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

Pesticide Disposal:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, add water to fill 10 percent of bottle. Agitate vigorously and spray onto labeled use site. Repeat this rinsing procedure two more times.

Completely empty container. Dispose of container at an approved waste disposal facility or by incineration if allowed by state and local authorities. Do not use container in connection with food, feed, or drinking water.

Spills:

If the container is leaking or material is spilled, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away.

You may contact Chemtrec at 800-424-9300 for decontamination procedures or any other assistance that may be necessary.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer HealthCare LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER HEALTHCARE LLC MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer HealthCare LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER HEALTHCARE LLC DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER HEALTHCARE LLC'S ELECTION, THE REPLACEMENT OF PRODUCT.

Bayer, the Bayer Cross and QuickBayt are registered trademarks of Bayer.
Bitrex is a registered trademark of Johnson Matthey Public Limited Company.



Manufactured by
Bayer HealthCare

Bayer HealthCare LLC
Animal Health Division
P.O. Box 390
Shawnee Mission, KS 66201, USA

81147923, R.0

BR021609