



# Tempo<sup>®</sup> SC Ultra Pest Control Concentrate



**Tempo<sup>®</sup> SC Ultra is now available in a small 32 mL size.**

**Tempo SC Ultra is packaged and priced for individual applications.**

**Broad-spectrum.**

- 5th-generation pyrethroid (beta-cyfluthrin) provides strong insecticidal activity in a concentrated liquid formula.
- Kills over 60 pests, including flies, ants and spiders.
- For use in and around barns, stables and other animal quarters.

**Just the right size.**

- One 32 mL bottle makes 2 gal of spray, enough to treat 2,000 sq ft.
- Display carton contains 12 individually packaged bottles.
- Individual packages can be presented in a display carton or on hooks, or be free-standing.

**Flexible and user-friendly.**

- Apply indoors as a spot, crack or crevice spray, or outdoors as a perimeter treatment.
- Active ingredient distributes evenly during application and stays in suspension without reagitation.
- Controls crawling, flying and wood-infesting pests.



**Tempo<sup>®</sup> SC Ultra**  
Pest Control Concentrate

# Tempo<sup>®</sup> SC Ultra Pest Control Concentrate

Kills crawling, flying, and wood-infesting insects and protects barns, horse stables, and other animal quarters from invading pests. Just mix and spray to provide effective control.

- ✓ Kills over 60 pests, including flies, ants, and spiders.
- ✓ Makes 2 gallons of spray, enough to treat 2,000 sq ft.
- ✓ Use in and around barns, horse stables and animal quarters.

ACTIVE INGREDIENT: Beta-cyfluthrin [CAS No. 68359-37-5] .....11.8%  
OTHER INGREDIENTS: .....88.2%  
Net Contents: 1.082 fl. oz (32 ml) 100.0%

**STOP - Read the label before use.  
Keep out of reach of children.**

## CAUTION PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice</li> <li>• Have a person sip a glass of water if able to swallow</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor</li> <li>• Do not give anything by mouth to an unconscious person</li> </ul>
<b>IF ON SKIN</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing</li> <li>• Rinse skin immediately with plenty of water for 15 – 20 minutes</li> <li>• Call a poison control center or doctor for treatment advice</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible</li> <li>• Call a poison control center or doctor for further treatment advice</li> </ul>
<b>HOTLINE NUMBER:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment call 800-422-9874. For product information call 800-633-3796.	
<b>Note to Physician:</b> No specific antidote is available. Treat the patient symptomatically.	

## ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment wash waters. Apply this product only as specified on this label. This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply Tempo SC Ultra Pest Control Concentrate or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

Tempo <sup>®</sup> SC Ultra Pest Control Concentrate	
<ul style="list-style-type: none"> <li>• For indoor and outdoor use</li> <li>• Use in and around animal facilities</li> <li>• Kills over 60 pests</li> <li>• Makes 2 gallons of spray, enough to treat 2,000 sq ft.</li> <li>• Non-staining</li> </ul>	
<b>Kills</b>	Ants, Fleas, Flies, Ticks, Spiders, and over 60 other listed pests
<b>Where to Use</b>	Indoors as a spot, crack, or crevice treatment or outdoors as perimeter treatment in and around barns, horse stables and other animal quarters. Reapplication can be made at 10-day intervals, if necessary.
<b>When to Use</b>	To control and prevent insect infestations
<b>How Much To Use</b>	Dilute contents of container (32 ml) in two gallons of water

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**IMPORTANT:** Read these entire Directions for Use and Conditions of Sale before using Tempo SC Ultra Pest Control Concentrate.

**CONDITIONS OF SALE:** The directions on this label were determined through research to be appropriate for the correct use of this product. This product has been tested under different environmental conditions both indoors and outdoors similar to those that are ordinary and customary where the product is to be used. Insufficient control of pests or plant injury may result from the occurrence of extraordinary or unusual conditions, or from failure to follow label directions. In addition, failure to follow label directions may cause injury to animals, man, and damage to the environment. To the extent permitted by applicable law, Bayer offers, and the buyer accepts and uses, this product subject to the conditions that extraordinary or unusual environmental conditions, or failure to follow label directions are beyond the control of Bayer and are, therefore, the responsibility of the buyer.

## GENERAL INFORMATION

Tempo SC Ultra Pest Control Concentrate will provide effective control of the pests listed on this label. Tempo SC Ultra Pest Concentrate is intended for use as a general surface, spot, or crack and crevice spray in and around buildings and structures and their immediate surroundings. Permitted areas of use include, animal runs, animal holding areas, animal quarters, boarding kennels, chicken houses (when birds are not present), dairy barns, feed lots, horse stables, kennels, livestock barns and holding facilities, storage areas, storage rooms, trailers, turkey houses, utility rooms, and rooms in animal facilities.

Not for use in and around residential areas. Do not use in commercial food/feed handling establishments, restaurants or other areas where food is commercially prepared, processed or served. Do not use food utensils such as teaspoons or measuring cup for food purposes after use with insecticides.

Cleaning of all areas prior to use of Tempo SC Ultra Pest Control Concentrate will increase levels of control. Any foodstuffs infested with pests should be removed and destroyed.

## MIXING INSTRUCTIONS

Tempo SC Ultra Pest Control Concentrate is intended to be mixed with water and applied with pump style sprayer or trigger spray bottle. Add the appropriate amount of product while filling the sprayer with water, and shake to mix. Use 1 Tablespoon (16 milliliters) of Tempo SC Ultra Pest Control Concentrate per 1,000 ft<sup>2</sup> in 1 gallon of water to adequately cover the area being treated, but which will not allow dripping or run off to occur. Diluted spray mixture can be stored overnight and applied the following day; however, mixture should be shaken prior to use to thoroughly mix the product.

Tempo SC Ultra Pest Control Concentrate will not stain or damage painted or varnished surfaces, plastics, fabrics or other surfaces where water applied alone causes no damage.

## INDOOR PEST CONTROL

For use in animal housing, including, livestock barns, horse stables, and other animal quarters to control crawling and flying insect pests. Apply diluted spray as a general surface and/or a crack and crevice spray. Control will

be enhanced when facilities are cleaned and interior applications are supplemented with exterior perimeter treatments. Do not make interior applications of Tempo SC Ultra Pest Control Concentrate in areas of facility where animals are present. Do not make applications to animal feedstuffs or watering equipment.

## GENERAL SURFACE APPLICATIONS

For general pests, occasional invaders, and flying insects apply the diluted spray evenly to walls, floors, between, behind and beneath equipment, appliances, around floor drains, window and door frames, and on the underside of shelves, drawers and in similar areas. Applications may be made to floor surfaces along walls and around air ducts; however do not treat the entire area or floor or floor coverings. Pay particular attention to treating entry points and harborage areas. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas. It is recommended to wear safety glasses, goggles or face shield and a respirator when making a general surface treatment to overhead areas. For control of flying insects, make application to surfaces or directly where the pest collects or rests. Do not apply as a fog.

## SPOT, CRACK OR CREVICE APPLICATIONS

Use low spray pressure and pin-point application in specific areas such as: cracks and crevices in or behind baseboards, in floors, walls, expansion joints, areas around water and sewer pipes, wall voids, or voids in table legs and equipment where pests can hide, storage areas, closets, around water pipes, doors and windows, behind and under cabinets, sinks, and other equipment, the underside of shelves, drawers, as well as other similar areas. Do not apply where electrical short circuits could occur. Use a dust or dry bait in these areas.

## OUTDOOR PEST CONTROL

For control of flying insects outdoors: Make applications to outside surfaces of barns, horse stables, animal confinement areas, animal quarters, buildings, feedlots, animal holding areas, storage areas, workshops, and other areas where these pests have been seen or found. Make pin stream applications directly to the nests of wasps, yellow-jackets and hornets. For perimeter pest control, make a general surface application to outside surfaces of barns, horse stables, animal confinement areas, animal quarters, buildings, feedlots, animal holding areas, storage areas, workshops, and other areas where pests have been seen or found. To help prevent infestations of buildings by occasional invaders, treat soil, mulch, turf or other substrates adjacent to buildings, the building foundation, walls, around doors and windows, soffit areas where these pests are active and may find entrance.

For control of the imported red fire ant, around the sites listed above, mix Tempo SC Ultra Pest Control Concentrate as directed above and thoroughly drench or inject the entire mound area. Treat all areas of the mound showing activity to provide contact control of the adult fire ants. Tempo SC Ultra Pest Control Concentrate will provide contact activity and control in the active mound.

## CONTROL OF WOOD INFESTING PESTS

Mix a diluted spray by mixing 1 Tablespoon (16 milliliters) of Tempo SC Ultra Pest Control Concentrate in one gallon of water.

For control of wood infesting beetles on structure surfaces apply diluted spray to thoroughly and evenly cover the area. If spraying overhead, cover all surfaces below the area being sprayed with plastic sheeting or other material which can be disposed of by placing in trash if contamination from dripping occurs. Sprayed surfaces should be avoided until the treated area is completely dry.

For control of above ground termites and wood-infesting beetles in localized areas, apply diluted spray to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable.

Such applications for termites are not a substitute for mechanical alteration, soil treatment or foundation treatment, but are merely a supplement. For active termite infestations, get a professional inspection.

PESTS CONTROLLED BY TEMPO <sup>®</sup> SC Ultra Pest Control Concentrate*			
INDOOR PESTS		OUTDOOR PESTS	WOOD INFESTING PESTS
<b>Crawling Pests</b>	<b>Flying Pests</b>		
Ants (except Pharaoh)	Clothes moths	Ants	Beetles
Bedbugs	Gnats	Bees	Ambrosia
Carpet beetles	Flies	Boxelder bugs	Deathwatch
Centipedes	Midges	Carpenter ants	False powderpost
Cockroaches	Mosquitoes	Cecidflies	Old house borer
American	Moths	Centipedes	Powderpost
Asian	Wasps	Cluster flies	Carpenter ants
German	Yellow-jackets	Cockroaches	Carpenter bees
Oriental		Crickets	Termites
brown-banded		Elm leaf beetle	Subterranean**
smoky brown		Fire ants	Formosan
Crickets		Firebrats	Drywood
Darkling beetles		Flies	
Earwigs		Fleas	
Firebrats		Gnats	
Millipedes		Ground beetles	
Pillbugs		Hornets	
Silverfish		Imported red fire ants	
Sowbugs		Midges	
Spiders		Millipedes	
		Mosquitoes	
		Moths	
		Scorpions	
		Silverfish	
		Spiders	
		Ticks	
		Wasps	
		Yellow-jackets	

\* For Outdoor and Indoor Pest Control use 1 Tablespoon (16 milliliters) of Tempo SC Ultra Pest Control Concentrate per 1000 square feet in one gallon of water.

\*\* Only for control of above-ground, winged forms of subterranean termites.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original containers and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away. For transportation emergencies contact Chemtrec at 800-424-9300.

**Pesticide Disposal:** Pesticide wastes are hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Residue Removal:** Triple rinse as follows: Empty the remaining contents into sprayer or spray bottle and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into sprayer or spray bottle for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Container Disposal:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

EPA Reg. No. 11556-145

EPA Est. No. 84926-IL-1

© 2008 Bayer HealthCare LLC

Bayer, the Bayer Cross and Tempo are trademarks of Bayer.

BR040908

80694254, R.0

13945



Bayer HealthCare LLC  
Animal Health Division  
P.O. Box 390  
Shawnee Mission, KS 66201