

FOR BROAD-SPECTRUM PEST CONTROL

Pick up the **Tempo**[®] family of products.



Tempo 1% Dust Insecticide
Tempo 20 WP Insecticide
Tempo SC Ultra Premise Spray



Tempo for VERSATILITY:

Registered for use in and around livestock buildings, including:

- Dairy barns
- Poultry houses
- Swine houses
- Milking parlors
- Hospital pens
- Clinics
- Chutes
- Horse stables

Works on crawling and flying pests.

Tempo for EFFECTIVENESS:

Fast-acting, working primarily as a contact insecticide with good activity through ingestion.

Efficacious at lower rates than other insecticides; 4–6 times more effective than 3rd-generation pyrethroid insecticides.

Stabilized 4th-generation pyrethroid with Cyfluthrin/
Beta-cyfluthrin active ingredients:

- Acts on both contact and ingestion
- Controls insects resistant to organophosphates and carbamates

GENERAL INSTRUCTIONS:

Do not use Tempo with animals present.

Avoid contact with skin and eyes. Wear protective clothing and an approved respirator when applying Tempo.

In cases of severe infestations, Tempo can be reapplied every 7–10 days.

Do not mix Tempo products with incompatible disinfectants. Complete a jar compatibility test before mixing.

Note: For best results, pre-acidify mix tank water to a pH of 5.0 or less prior to mixing with Tempo 20 WP and Tempo SC Ultra formulations.*

ALWAYS READ AND FOLLOW LABEL DIRECTIONS BEFORE USE



Tempo®

20 WP Insecticide



Superior formulation for rough surfaces and areas containing large amounts of dust and organic debris

FACTS:

20% cyfluthrin wettable powder

Can be tank-mixed with other pesticides registered for similar uses

Packaged in 420-gram jar

INSTRUCTIONS:

Use tanks with agitating mixers, and frequently agitate during application.

For general surface applications and/or longer residual for severe infestations to end walls, sidewalls, rafters and ceilings, and for spot treatment of areas under equipment (such as feed lines or feed troughs), mix one jar of Tempo 20 WP in 21 gallons of water to create a 0.1% solution.

COVERAGE:

10 to 20 grams treats 1,000 square feet.

Tempo®

1% Dust Insecticide



The easy-to-apply dust that requires no mixing equipment and remains effective in cracks and crevices the longest

FACTS:

Provides effective pest control in areas of high humidity

Little to no odor

Not corrosive

Available in sleeves of 4 inside 10-lb bucket

INSTRUCTIONS:

Apply dust lightly and uniformly to all infested areas using hand-held, vehicle-mounted or backpack dusters.

Close all doors, curtains, vents or other openings and turn off fans prior to dust applications.

Allow 48 hours between premise dust treatment and reintroduction of animals.

COVERAGE:

0.5 to 1 lb treats 1,000 square feet.

Tempo®

SC Ultra Premise Spray



The strongest insecticidal activity in a stable, suspension-concentrated liquid formula

FACTS:

5th-generation pyrethroid (with its beta-cyfluthrin active ingredient) provides the most potent insecticidal activity

Allows for more even disbursement during application

Active ingredient stays in solution without reagitation

Packaged in 240-mL and 900-mL tip-and-pour plastic bottle

INSTRUCTIONS:

For general surface applications and/or longer residual for severe infestations to end walls, sidewalls, rafters and ceilings, mix one 240-mL bottle of Tempo SC Ultra per 45 gallons of water (for unpainted particle board or cement block).

For spot treatment of areas under equipment (such as feed lines or feed troughs), mix one 240-mL bottle with 15 gallons of water.

COVERAGE:

8 to 16 mL treats 1,000 square feet.

PESTS CONTROLLED BY TEMPO® SC ULTRA PREMISE SPRAY

INDOOR PREMISE PESTS			OUTDOOR PERIMETER PESTS		WOOD-INFESTING PESTS
Crawling Pests	Flying Pests	Pantry & Stored-Product Pests	Ants	Gnats	Beetles
Ants (except Pharaoh)	Clothes moths	Beetles (exposed adults & immature stages)	Bees	Ground beetles	– Ambrosia
Bed bugs	Flies	– Cadelle	Boxelder bugs	Hornets	– Deathwatch
Carpet beetles	Gnats	– Cigarette	Carpenter ants	Imported red fire ants	– False powderpost
Centipedes	Midges	– Confused flour	Cecidflies	Midges	– Old house borer
Cockroaches	Mosquitos	– Dermestid	Centipedes	Millipedes	– Powderpost
– American	Moths	– Drugstore	Cluster flies	Mosquitoes	Carpenter ants
– Asian	Wasps	– Granary weevil	Cockroaches	Moths	Carpenter bees
– German	Yellow-jackets	– Hide	Crickets	Scorpions	Termites
– Oriental		– Larder	Elm leaf beetle	Silverfish	– Subterranean
– brownbanded		– Leather	Fire ants	Spiders	– Formosan
– smokybrown		– Lesser grain borer	Firebrats	Ticks	– Drywood
Crickets		– Lesser mealworm	Flies	Wasps	
Darkling beetles		– Red flour	Fleas	Yellow-jackets	
Earwigs		– Rice weevil			
Firebrats		– Sawtoothed grain			
Millipedes		– Warehouse			
Pillbugs		Moths			
Silverfish		– Indianmeal			
Sowbugs		– Mediterranean flour			
Spiders					

Tempo®

SC Ultra Premise Spray

For broad-spectrum control of crawling, flying, and wood-infesting insect pests in and around animal housing, warehouses, and processing and packing plants.

Shake well before using.

ACTIVE INGREDIENT:

Beta-cyfluthrin; Cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate 11.8%

OTHER INGREDIENTS 88.2%
100.0%

Contains 1 lb cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropanecarboxylate per gallon.

STOP - Read the label before use. Keep out of reach of children.

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove contaminated clothing and wash before reuse.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center for treatment advice. **If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product container with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or when disposing of equipment wash waters. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply Tempo® SC Ultra insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire Directions for Use and Conditions of Sale before using Tempo® SC Ultra Insecticide.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

GENERAL INFORMATION FOR STRUCTURAL PEST CONTROL

Tempo® SC Ultra Insecticide is a liquid formulation containing 1 lb active ingredient per gallon, which will provide effective knockdown and residual control of the pests listed in this label.

Tempo® SC Ultra is intended for use as a general surface, spot, mist, or crack and crevice application in and around buildings and structures and their immediate surroundings. Permitted areas of use include, **but are not limited to**, dairies and dairy product processing plants, grain mills, granaries, greenhouses (structures only), industrial buildings, kitchens, laboratories, livestock housing structures, pet kennels, manufacturing establishments, poultry houses, meat, poultry and egg processing and packing plants, meat and produce canneries, stores, warehouses and similar structures.

Tempo® SC Ultra is intended to be mixed with water and applied with hand pressurized, power operated sprayers or foam generating equipment. Spray pressure should not exceed 50 PSI at the nozzle tip when used indoors. Add the appropriate amount of product when filling sprayer tank with water, shake or agitate as necessary to mix. Diluted spray mixture can be stored overnight and applied the following day; however, mixture should be agitated prior to application to prevent uneven distribution of product. Before leaving spray mixture unagitated for an extended period of time, release tank pressure and elevate nozzle and hose above tank outlet, activate trigger and drain all dilution from hose back into spray tank. Re-tighten spray tank lid and do not re-pressurize until starting spray again.

Tempo® SC Ultra will not stain or cause damage to any painted or varnished household surface, plastic, fabric or other surface where water applied alone causes no damage.

TANK MIXING: The user, at his discretion, can tank mix pesticides currently registered for similar use patterns, unless the product labels specifically prohibit such use. See section **Extended Cockroach Control with Insect Growth Regulators**. It is always recommended that a small jar compatibility test using proper proportions of chemicals and water be run to check for physical compatibility prior to tank mixing. Tempo® SC Ultra is not compatible with disinfectants and should not be tank mixed with them. Tempo® SC Ultra and disinfectants should be applied separately as per label directions.

GENERAL SURFACE APPLICATION

Use 8 milliliters of Tempo® SC Ultra per 1000 sq ft in sufficient water (under conditions of severe pest infestation or when quicker knock-down and/or residual control is needed, use 16 milliliters of Tempo® SC Ultra per 1000 sq ft) to adequately cover the area being treated, but which will not allow dripping or run-off to occur (see Note 1). Use a low pressure system with a fan type nozzle to uniformly apply the dilution. Applications can be made to walls, floors, ceilings, in and around cupboards, between, behind and beneath equipment, appliances, around floor drains, window and door frames, and on the underside of shelves, drawers and in similar areas. Applications may be made to floor surfaces along walls and around air ducts, however, do not treat entire area of floor or floor coverings. All food processing

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Remove from premises or tightly cover fish tanks and disconnect aerators when applying indoors where such containers are present. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or residues on crops or weeds. Do not apply TEMPO 20 WP Insecticide or allow it to drift onto crops or weeds on which bees are actively foraging. Additional information may be obtained by consulting your Cooperative Extension Service. See leaflet accompanying this container for complete labeling including Directions for Use and Storage and Disposal.

PRO FACTS

- **Quick Knockdown & Residual Control**
- **Low Odor**
- **Non-staining**

TEMPO 20 WP	• Wettable powder containing 20% cyfluthrin
CONTROLS	• Indoor Pests • Outdoor/ Perimeter Pests <i>For list of specific insects, see inside.</i>
WHERE TO APPLY	Apply where pests have been seen or found, or can find shelter. • In and around buildings and structures
COVERAGE	• Makes up to 42 gallons of finished spray solution for indoors.(420 g size) • Treats up to 42,000 sq ft. outdoors. (420 g size)
MIXING INFORMATION	• Mix the appropriate amount of TEMPO 20 WP with water. • May tank mix with other pesticides currently registered for similar uses, unless specifically prohibited.
HOW TO APPLY	• General surface spot, mist, or crack & crevice application.
RE-APPLY	• Every 7 to 10 days, if needed.
RE-ENTRY	• Avoid contact with treated surfaces until dry. • People and pets may re-enter after the treated area is dry.
QUESTIONS	For questions or comments, call toll-free 1-800-633-3796 www.bayer-ah.com

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read entire label and Conditions of Sale before using TEMPO 20 WP Insecticide.

Do not formulate this product into other end-use products.

STRUCTURAL PESTS

FOR RESIDUAL PEST CONTROL IN AND AROUND BUILDINGS, STRUCTURES AND FOOD/FEED HANDLING ESTABLISHMENTS. Use TEMPO 20 WP as a general surface, spot, mist, or crack and crevice treatment in and around buildings and structures. This includes factories, warehouses, and food processing facilities. For a list of common facilities where this product may be used, see Appendix.

INDOOR PESTS

General Household Pests

Crawling Pests

- Ants (except Pharaoh)
- Bedbugs
- Carpet beetles
- Centipedes
- Clothes moths
- Cluster fly

- Pillbugs
- Silverfish
- Sowbugs
- Spiders

Flying Pests

- Flies
- Gnats
- Hornets
- Midges
- Moths
- Wasps
- Yellow jackets

Beetles

(exposed adults & immature stages)

- Cadelle
- Cigarette
- Confused flour
- Dermestid
- Drugstore

- Granary weevil
- Hide
- Larder
- Leather
- Lesser grain borer
- Lesser mealworm
- Merchant grain

- Mealworm
- Red flour
- Rice weevil
- Sawtoothed grain
- Warehouse

- Moths
- Indian meal
- Mediterranean flour

WHERE TO APPLY	Apply where pests have been seen or found, or can find shelter. Use a general surface, spot, mist, or crack and crevice application.
USE RATE	0.05%—10 g (2 scoops)/gal water 0.1%—20 g (4 scoops)/gal water • Use high rate for severe infestations, faster knockdown, and longer residual. • Always use the high rate for bees, wasps, yellow jackets and hornets.
HOW TO APPLY (Also see APPLICATION FOR INDOOR PESTS)	• Remove or tightly cover fish tanks and disconnect aerators during application. • Remove pets during application and until treated surfaces are completely dry. • Avoid contact with treated surfaces until dry. • Wear safety glasses, goggles, or face shield and a dust/mist respirator when treating overhead areas. General Surface Applications • Use a low-pressure system (do not exceed 50 psi at nozzle tip), with a fan-type nozzle to apply the dilution uniformly. Spot, Mist, or Crack and Crevice Applications • Apply directly into voids, cracks and crevices using spray equipment. • Use a low-pressure system (do not exceed 50 psi at nozzle tip), with a pinpoint or variable pattern nozzle or applicator equipment that delivers low volume treatments, such as the Micro-Injector® or Actisol®.
WHEN TO APPLY	• When pests are seen or found • Re-apply every 7 to 10-days if needed.
MIXING INFORMATION	• Add product when filling spray tank with water; shake or agitate mixture. • Diluted spray mixture can be stored overnight. Agitate before using. Can be tank mixed with other pesticides currently registered for similar uses. To be sure of physical compatibility, first prepare a sample in a small jar using the proper proportions of water and products. Follow the most restrictive of labeling directions and precautions. Do not exceed label rates.
RESTRICTIONS	• Do not treat entire area of floor or floor coverings. • Do not apply where electrical short circuits can occur. Use a dust or dry bait in these areas. • Do not apply to furniture or upholstery where prolonged contact by humans will occur. • Do not apply as a space spray.
ADDITIONAL DIRECTIONS FOR FOOD AND FEED HANDLING ESTABLISHMENTS	

WHERE TO APPLY	Applications may be made to both Food/Feed and non-food areas of food and feed handling establishments. Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared, or served. Includes areas for: • receiving, storing and packing food • preparing food • edible waste storage • enclosed processing systems (mills, dairies, edible oils, syrups) of food. Serving areas where food is exposed and the facility is in operation are also considered food areas.
USE RATE	• Apply as a general surface spot, mist, or crack & crevice application. 0.05%—10 g (2 scoops)/gal water 0.1%—20 g (4 scoops)/gal water • Use high rate for severe infestations, faster knockdown, and longer residual.

HOW TO APPLY

(Also see INDOOR PESTS)

General Surface Applications

Cover or remove all food processing and handling equipment before treatment.

After applying where food is commercially prepared or processed, wash all handling and processing equipment, benches, shelving, and other surfaces which food will contact. Rinse thoroughly with clean, fresh water.

Do not apply to surfaces in food-handling areas when facility is in use or when foods are exposed.

Do not apply directly to food.

Spot or Crack & Crevice Applications

Spot or crack and crevice application is permitted while facility is in use. Cover or remove food from area being treated.

Do not apply directly to food or food handling surfaces.

APPLICATION FOR INDOOR PESTS

GENERAL HOUSEHOLD PESTS

- Cockroaches, Crickets, Silverfish, Spiders, and others

- Use a coarse, low-pressure spray.
- Treat entry points such as around water pipes, doors, windows, and eaves. Treat areas where pests normally feed or hide such as baseboards, corners, around water pipes, under refrigerators, cabinets, sinks, and stoves, being sure spray cracks & crevices. Spot treat floor or rugs beneath furniture, in closets, and storage areas, but do not apply to entire floor area.
- **Do not** use as a "space spray".
- **IGR Note:** This product can be tank mixed with an insect growth regulator (IGR) for extended cockroach control.

ANTS

- Apply to ant trails on walls, tile, baseboards, around pipes, under cabinets, and other places where ants are active.
- Treat entry points such as windows and doors and the outdoor perimeter.
- For best results, treat nests when possible.

PANTRY AND STORED PRODUCT PESTS

- For best results, clean area before treating. Remove and destroy infested food. Discard food accidentally contaminated with spray.
- For cupboards, shelving, or food storage areas: Remove utensils, exposed food, shelf paper, and other items before spraying. When dried, cover shelves with clean paper before replacing items.

PESTS IN LIVESTOCK HOUSING STRUCTURES
including poultry houses & pet kennels

- Apply as a general surface and/or a crack and crevice spray.
- For best results, clean facilities before applying and treat outdoor perimeter areas.
- **Do not** apply indoors where animals other than cattle or horses are present.
- **Do not** apply to animal feedstuffs or watering equipment.

FLYING INSECTS

- Apply to surfaces where pests collect or rest.

OUTDOOR/PERIMETER PESTS

Ants	Cockroaches	Firebrats	Hornets	Scorpions	Wasps
Bees	Cluster fly	Fleas	Midges	Silverfish	Yellow jackets
Boxelder bugs	Crickets	Flies	Millipedes	Sowbugs	
Carpenter ants	Elm leaf beetle	Gnats	Moths	Spiders	
Centipedes	Fire ants	Ground beetles		Pillbugs	Ticks

WHERE TO APPLY

- Apply where pests have been seen or found, or can find shelter.
- Apply to buildings, porches, patios, garages, fences and other areas where pests are found.
- To prevent infestation, treat the building foundation, walls, and around doors, windows, and soffits where pests are active or may enter or hide; also treat the soil, turf, or other substrates adjacent to buildings

USE RATE

- 10 to 20 g (2 to 4 scoops) / 1000 sq ft See Table 1 for mixing directions.
- For fire ants and flying pests
0.05% —10 g (2 scoops) /gal water
0.1% — 20 g (4 scoops) /gal water
- Always use the high rate for fire ants, bees, wasps, yellow jackets and hornets.
- Use high rate for severe infestations, faster knock down, and longer residual.

HOW TO APPLY
(Also see

- **Perimeter Treatment:** To help prevent infestation of structures, treat area adjacent buildings in a band up to 10 feet wide. Also treat the building foundation to a height of 2 to 3 feet in areas where pests are active or can find entrance.
 - Apply just enough dilution to adequately cover the area without excessive dripping or runoff.
- Volume can vary depending on the surface type treated. For example, mulch requires more spray than bare soil. If mulch is heavy, rake it back to expose soil before treating.
- Avoid contact with treated surfaces until dry.

WHEN TO APPLY

- When pests are seen or found
- Re-apply every 7 to 10-days if needed.

MIXING INFORMATION

- Add product when filling spray tank with water; shake or agitate mixture.
- Diluted spray mixture can be stored overnight. Agitate before using.
- Can be tank mixed with other pesticides currently registered for similar uses. To be sure of physical compatibility, first prepare a sample in a small jar using the proper proportions of water and products. Follow the most restrictive of labeling directions and precautions. Do not exceed label rates.

TABLE 1: Mixing Directions for Outdoor/Perimeter Applications

Application Volume: Gallons/1,000 sq. ft.	Use Rate: Grams/1,000 sq. ft.	Scoops* of TEMPO 20 WP Diluted to These Volumes of Finished Spray			
		3 gal.	10 gal.	15 gal.	25 gal.
2	10	3	10	15	25
	20	6	20	30	50
5	10	1	4	6	10
	20	2	8	12	20
10	10	—	2	3	5
	20	—	4	6	10

*1 level scoop = 5 grams

FORMULA FOR DETERMINING THE PERCENT CONCENTRATION IN THE FINISHED SPRAY

$$\frac{\text{(Scoops of TEMPO 20 WP)}}{\text{(Gallons of finished spray)}} \times 37.85 = \text{Percent Concentration}$$

APPLICATION FOR OUTDOOR/PERIMETER PESTS

Ants	Locate and treat ant nests, if possible. Follow trails of foraging workers, check under objects near the structure, and inspect other likely nest sites. If ants are nesting in the soil, thoroughly drench or inject the nest and surrounding soil. Treat ant trails around windows and doors, along sidewalks, or other areas where seen. Apply a perimeter treatment as described above.
Bees, Wasps, Yellow Jackets, Hornets	Use Rate: 0.1% concentration. Make pin stream applications directly to the nests. To help avoid stings treat nests in the evening when the insects are inside the nest. For best results, check nests within one or two days to ensure complete kill. Remove and destroy treated nests to prevent emergence of newly hatched insects, secondary pest problems, and possible staining from honey. If removal is not possible, retreat nests if necessary.

Boxelder Bugs	Apply directly to boxelder bugs that have congregated around doors and windows and similar areas where they may enter structures. Apply perimeter treatment as described above. For best results, treat infested trees early in the season. Tree spraying may require ornamental license—check with your local regulator.
Centipedes, Millipedes, Pillbugs, Sowbugs	Apply perimeter treatment to reduce the number of insects from entering the structure. Thoroughly treat mulched areas. Reduce moisture areas from around the structure.
Fire Ants	Use Rate: 0.1% concentration Thoroughly drench or inject the entire mound. When using injection treatment, you must treat the bottom of the mound. Apply dilution using a downward and upward motion of the injector tool. For complete distribution, inject in several points throughout the mound. Treat new mounds as they appear. Do not use in pastures or areas of food crop production.
Fleas & Ticks	Make a perimeter treatment, concentrating on areas where pets rest or run. For ticks, treat likely harborage areas like along paths, tall grass and other vegetation. For best results, treat the entire area where ticks could be present. Begin treating in the spring. Ticks may be re-introduced from surrounding areas by host animals. Continue treating until frost to control both larvae and adult ticks.
Flies	Use Rate: 0.05% to 0.1% Apply low-volume mist or spray to outside surfaces of buildings, porches, patios, garages, fences, and other areas where flies rest. CLUSTER FLY: Treat in late summer or fall, just before flies begin looking for winter shelter. Spray siding, eaves, around windows and doors, window screens, and attics. To help reduce infestations, treat cracks and crevices, attics and other indoor winter hiding places. Note: Cluster flies may appear during winter warm spells, while snow is still on the ground.
Mosquitoes	Spray around landscape plants, turf and ground cover, under decks, around building foundations and other places where mosquitoes may rest.
Scorpions	Treat around piles of firewood and other areas where scorpions may hide. When possible, remove piles of materials that can harbor the pests. Apply a perimeter treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage:	<ul style="list-style-type: none"> Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Protect from freezing. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.
Pesticide Disposal:	<ul style="list-style-type: none"> Pesticide wastes are hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
Container Disposal:	<ul style="list-style-type: none"> Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment. Triple rinse, or equivalent. Dispose of container by offering for recycling or reconditioning. Or puncture and dispose of container in a sanitary landfill, or by incineration, or, if allowed by state and local authorities by burning. If burned, stay out of smoke.
Spills:	<p>If the container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material.</p> <ul style="list-style-type: none"> Do not walk through spilled material. Dispose of as directed for pesticides above. Keep unauthorized people away. You may contact the Bayer Emergency Response Team for decontamination procedures or any other assistance at 800-334-7577 or contact Chemtrec at 800-424-9300.

CONDITIONS OF SALE. Read Before Using.

THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

©2005 Bayer HealthCare LLC

APPENDIX

Use TEMPO 20 WP as a general surface, spot, mist, or crack and crevice treatment. Permitted areas of use in buildings, structures, and modes of transport include, *but are not limited to*:

- dairies and dairy product processing plants
- grain mills
- granaries
- livestock housing
- pet kennels
- poultry houses
- warehouses
- transportation vessels such as trucks, trailers, and rail cars

Net Contents 14.8 oz (420 Grams)

08711286-035999, R.6

EPA Reg. No. 432-1302-11556

BR080907

EPA Est. No. 33967-NJ-1



Manufactured for Bayer HealthCare LLC, Animal Health Division,
Shawnee Mission, Kansas 66201 U.S.A.
Bayer, Bayer Cross and Tempo are registered trademarks of Bayer.

Tempo®

1% Dust Insecticide

For Control Of Crawling And Flying Insect Pests
In Poultry And Livestock Facilities.

ACTIVE INGREDIENT:

Cyfluthrin, cyano(4-fluoro-3-phenoxyphenyl)methyl 3-(2,2-dichloroethyl)-2,2-dimethylcyclopropanecarboxylate1%

OTHER INGREDIENTS99%

Percent

By Weight

100%

STOP - Read The Label Before Use

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUCIÓN AL USUARIO: Si usted no puede leer o entender inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not get into eyes, on skin, or on clothing. Avoid breathing dust. If clothing becomes contaminated, remove and wash before reuse. Wash hands thoroughly with soap and warm water after handling. Keep out of reach of children. Do not contaminate feed or food.

SYMPTOMS OF POISONING: In case of poisoning, call physician or Poison Control Center immediately. Have patient lie down and keep quiet.

FIRST AID	
If in eyes:	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none"> Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on skin:	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
HOTLINE NUMBER: Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency medical treatment information call 1-877-258-2280. For product information call 1-800-633-3796.	
Note to Physician: No specific antidote is available. Treat the patient symptomatically.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or remove any fish or aquatic organism tank within structures being treated. Disconnect any air supply sources to such tanks before applications are made and reconnect after any possibility of contamination has passed. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Disposal: Pesticide wastes are hazardous. Improper disposal of excess pesticide is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Do not use container in connection with food, feed, or drinking water. Completely empty container into application equipment. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original containers and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material is spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Dispose of pesticide as directed above. In spill or leak incidents, keep unauthorized people away.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

IMPORTANT: Read these entire **Directions for Use** and **Conditions of Sale** before using TEMPO 1% Dust Insecticide.

Do not formulate this product into other end use products.

GENERAL INFORMATION

TEMPO 1% Dust Insecticide is a ready-to-use insecticide dust which provides effective knockdown and residual control of the pests listed on this label.

TEMPO 1% Dust Insecticide is for residual pest control in and around livestock and poultry housing facilities. Permitted areas of use include, **but are not limited to**, livestock housing structures, poultry houses, swine houses, cattle barns, horse stables and pet kennels.

Do not use in food/feed handling establishments, restaurants or other areas where food/feed is commercially prepared or processed.

INDOOR PESTS

Apply dust using hand or power dusters or other suitable equipment. Apply lightly and uniformly to infested areas. Pay particular attention to floors, walls and bedding, as well as, cracks and crevices; wall voids and around window and door frames. Apply 0.5 to 1 pound of TEMPO 1% Dust Insecticide per thousand square feet. Do not make applications of TEMPO 1% Dust when animals are present in area of facility to be treated. Do not apply directly to animal feedstuffs or watering equipment. Repeat treatments as necessary to maintain adequate control.

TEMPO 1% Dust Insecticide controls the following indoor pests: ants (except pharaoh), bedbugs, beetles (exposed adults & immature stages), fire ants, moths, spiders, wasps and yellow-jackets.

OUTDOOR PESTS

Wasps and Bees: It is advisable to treat wasp and bee nests in the evening when insects are less active and have returned to the nest. Wear protective clothing if deemed necessary to avoid stings. Using hand-held or power duster or other suitable means, with extension tubes if necessary, thoroughly dust nest and entrance and surrounding areas where insects alight. To treat nests in wall voids, drill a hole in the area, dust entrance and surrounding areas. For best results check nests carefully one or two days after treatment to ensure complete kill, then remove and destroy nest to prevent emergence of newly-hatched insects. If removal is not feasible, retreat the nest, if necessary.

CONDITIONS OF SALE: THE DIRECTIONS ON THIS LABEL WERE DETERMINED THROUGH RESEARCH TO BE APPROPRIATE FOR THE CORRECT USE OF THIS PRODUCT. THIS PRODUCT HAS BEEN TESTED UNDER DIFFERENT ENVIRONMENTAL CONDITIONS BOTH INDOORS AND OUTDOORS SIMILAR TO THOSE THAT ARE ORDINARY AND CUSTOMARY WHERE THE PRODUCT IS TO BE USED. INSUFFICIENT CONTROL OF PESTS OR PLANT INJURY MAY RESULT FROM THE OCCURRENCE OF EXTRAORDINARY OR UNUSUAL CONDITIONS, OR FROM FAILURE TO FOLLOW LABEL DIRECTIONS. IN ADDITION, FAILURE TO FOLLOW LABEL DIRECTIONS MAY CAUSE INJURY TO ANIMALS, MAN, AND DAMAGE TO THE ENVIRONMENT. BAYER OFFERS, AND THE BUYER ACCEPTS AND USES, THIS PRODUCT SUBJECT TO THE CONDITIONS THAT EXTRAORDINARY OR UNUSUAL ENVIRONMENTAL CONDITIONS, OR FAILURE TO FOLLOW LABEL DIRECTIONS ARE BEYOND THE CONTROL OF BAYER AND ARE, THEREFORE, THE RESPONSIBILITY OF THE BUYER.

EPA Reg. No. 11556-136

EPA Est. No. 432-TX-1

SUPPLIED: 4 x 10 Pounds (4.50 kg)

08711294-037599, R.3

10 Pounds (4.50 Kg)

08715575-80003750, R.3

BR080907



Bayer HealthCare LLC, Animal Health Division,
Shawnee Mission, Kansas 66201 U.S.A.

